

## Claims

1-17 (Cancelled).

18. (Currently Amended)

A method ~~Computer-readable medium~~ for identifying a most likely biological pathway of a set of interacting molecules, wherein said interacting molecules each have one or more conserved features, the method comprising using a computer to execute instructions in a ~~the computer readable medium comprising the instructions operable to perform the steps of:~~

- a. representing the set of interacting molecules as oriented network graph,  $G = \langle V, E \rangle$ , where the vertices,  $V$ , correspond to molecules, and the edges or connections  $E$ , correspond to interactions between molecules;
- b. assigning a probability  $P(\text{network})$  to each possible oriented network graph found from a fixed number of vertices  $V$ ; and
- c. selecting an oriented network graph as the most likely biological pathway based on the probabilities  $P(\text{network})$  assigned to all possible oriented network graphs, whereby the selected graph provides a molecular interaction network representation of the biological pathway,

wherein step (b) comprises:

(i) assigning an attraction probability  $p_{ij}$  to every pair of molecules  $i$  and  $j$  that the molecules are connected to each other by an oriented edge, and conversely a probability  $(1-p_{ij})$  that the molecules are not connected to each other by an oriented edge;

(ii) computing a probability  $P(E)$  that a network has a particular edge set  $E$  ( $\epsilon_{ij}$ );

(iii) sorting all the possible oriented network graphs into a finite number of bins, each corresponding to a particular network topology that corresponds to a particular distribution of edges coming into and out of each vertex in the particular network, so that each bin represents a collection of oriented network graphs characterized by the same network topology;

(iv) computing a probability  $P(\text{topology})$  for each bin; and

(v) computing  $P(\text{network})$  as the product of  $P(E)$  and  $P(\text{topology})$ .

19. (Currently Amended) The ~~computer-readable-medium~~ method of claim 18, wherein said attraction probabilities  $p_{ij}$  are dependent on the conserved features of each molecule and are determined by quantifying the occurrence frequency of said features immediately upstream or downstream of each other within the known networks.

20. (Cancelled)

21. (Currently Amended) The ~~computer-readable-medium~~ method of claim 18, wherein said attraction probabilities  $p_{ij}$  are determined by quantifying the number of times conserved features in every pair of molecules  $i$  and  $j$  are seen in interaction with each other in known networks.

22. (Cancelled).

23. (Cancelled).

24. (Cancelled)

25. (Currently Amended) The ~~computer-readable-medium~~ method of claim 18, further comprising instructions operable to perform the step of obtaining a posterior probability of each possible oriented network graph, and wherein said selected oriented network graph corresponds to the graph having the highest posterior probability.

26. (Currently Amended) The ~~computer-readable-medium~~ method of claim 25, wherein said posterior probability is determined by using Markov Chain Monte Carlo techniques.
27. (Currently Amended) The ~~computer-readable-medium~~ method of claim 18, wherein said interacting molecules are proteins and said conserved features are protein domains or motifs.
28. (Currently Amended) The ~~computer-readable-medium~~ method of claim 27, wherein said conserved features are nucleic acid motifs.
29. (Currently Amended) The ~~computer-readable-medium~~ method of claim 27, wherein said attraction probabilities  $p_{ij}$  are determined by quantifying the occurrence frequency of said conserved features at immediately upstream or downstream of each other within the known networks; and said attraction probability  $p_{ij}$  of each pair of reaction molecules is determined by quantifying the number of times conserved features in every pair of molecules  $i$  and  $j$  are seen in interaction with each other in known networks.
30. (Cancelled)
31. (Currently Amended) The ~~computer-readable-medium~~ method of claim 27, wherein said probability  $P(\text{topology})$  of each possible network is a product of incoming edge distribution probability and outgoing edge distribution probability within said each possible network; and further comprising step of obtaining posterior probability of each network, wherein said molecular interaction network having the highest network probability is the network having the highest posterior probability, wherein said highest posterior probability is determined by Markov Chain Monte Carlo techniques.
32. (Currently Amended) The ~~computer-readable-medium~~ method of claim 31, wherein said attraction probabilities of said features are determined by quantifying the number of times conserved features in every pair of molecules  $i$  and  $j$  are seen in interaction with each other within the known networks.

33. (Currently Amended) The ~~computer-readable-medium~~ method of claim 31, wherein said attraction probabilities of said features are determined by quantifying the number of times conserved features in every pair of molecules i and j are seen in interaction with each other within the known networks.
34. (Cancelled)
35. (Cancelled)
36. (Currently Amended) A ~~computer-readable-medium~~ method for identifying a molecular interaction network representation for a set of interacting molecules within a known biological system, wherein each of said interacting molecules have one or more conserved features, the method comprising using a computer to execute instructions in a ~~the computer readable medium comprising instructions operable to~~ perform the steps of:
- a. determining attraction probabilities between pairs of molecules of the set of interacting molecules based on known molecular interaction data wherein said attraction probabilities are determined by quantifying the occurrence frequency of said conserved features of said pair of molecules at immediately upstream or downstream of each other within the known biological system networks;
  - b. determining an edge probability  $P(E)$  for ~~of~~ each possible molecular interaction network of the set of interacting molecules, based on the determined attraction probabilities of each pair of interacting ~~reaction~~ molecules within said each possible molecular interaction network;
  - c. determining a topology probability  $P(\text{topology})$  of said each possible molecular interaction network based on the network topology of said ~~each~~ network, said topology probability being a product of an incoming edge distribution probability and an outgoing edge distribution probability within said each possible molecular interaction network.

- d. determining a network probability of said each possible molecular interaction network as a product of said edge probability  $P(E)$  and said topology probability  $P(\text{topology})$ ;
- e. determining a posterior probability of said each possible molecular interaction network using equation 10; and
- f. identifying the possible molecular interaction network having the highest posterior probability as said molecular interaction network representation for the set of interacting molecules.

37. (Currently Amended) The ~~computer-readable medium~~ method of claim 36, wherein said molecule[s] is a protein and said conserved features are protein domain or motif.

38. (Currently Amended) A ~~computer-readable medium~~ method for identifying a molecular interaction of a molecule within a biological network of interacting molecules, said interacting molecules each having one or more conserved features, the method comprising using a computer to execute instructions in a ~~the computer readable medium comprising instructions operable~~ to perform the steps of:

- a. identifying a conserved feature of said molecule
- b. determining attraction probabilities between the conserved features of said molecule and other interacting molecules based on known molecular interaction data of the biological network;
- c. determining molecular interaction probabilities of said molecule with each of the other interacting molecules based on the attraction probabilities; and
- d. identifying the molecular interaction of said molecule with a one of the other interacting molecules, corresponding to the highest of the determined molecular interaction probabilities.

39. (Currently Amended) The ~~computer-readable medium~~ method of claim 38, wherein said attraction probabilities of said features are determined by using equation 6 or 14.
40. (Currently Amended) The ~~computer-readable medium~~ method of claim 38, wherein said attraction probabilities of said features are determined by using equation 14.
41. (Currently Amended) The ~~computer-readable medium~~ method of claim 38, wherein said molecular probability of the molecular interaction between said two molecules is identified by using equations 5 or 17.
42. (Currently Amended) The ~~computer-readable medium~~ method of claim 38, wherein said likelihood of molecular interactions between said interacting molecules is determined by using equations 17.
43. (Currently Amended) The ~~computer-readable medium~~ method of claim 38, wherein said molecules are protein, and said conserved features are protein domain or motif.
44. (Previously Presented) A screening method for identification of a compound capable of modifying the interaction between at least two molecules in a biology network comprising
- identifying an interaction between said at least two molecules using the method of claim 38;
  - introducing a test compound in the biology network, the test compound contacting said at least two molecules;
  - comparing the identified interaction of the molecules in the presence of the test compound with the identified interaction in the absence of the test compound;
- wherein a difference in the identified interaction of the molecules in the presence of the test compound as compared to the interaction in the absence of a test compound indicates identification of a compound capable of modifying the interaction between molecules.

45. (Previously Presented) The screening method of claim 44, wherein said molecules are proteins.